

Dynamics of F/D networks: the rôle of bound states

Mairi Sakellariadou¹ and Horace Stoica²

¹ Department of Physics, King's College London, London WC2R 2LS, U.K.

E-mail: Mairi.Sakellariadou@kcl.ac.uk

² Blackett Laboratory, Imperial College London, SW7 2AZ U.K.

E-mail: mairi.sakellariadou@kcl.ac.uk, f.stoica@imperial.ac.uk

Abstract.

We study, via numerical experiments, the rôle of bound states in the evolution of cosmic superstring networks, being composed by p F-strings, q D-strings and (p, q) bound states. We find robust evidence for scaling of all three components of the network, independently of initial conditions. The novelty of our numerical approach consists of having control over the initial abundance of bound states. This indeed allows us to identify the effect of bound states on the evolution of the network. Our studies also clearly show the existence of an additional energy loss mechanism, resulting to a lower overall string network energy, and thus scaling of the network. This new mechanism consists of the formation of bound states with an increasing length.

PACS numbers: 11.27.+d, 98.80.Cq

Submitted to: *JCAP*

1. Introduction

Cosmic superstrings can be formed as a result of brane annihilations in the context of brane-world scenario [1, 2, 3]. In the brane-world models, brane inflation takes place [4] while two branes move towards each other, and their annihilation releases the brane tension energy that heats up the universe to start the hot big bang era. Typically, strings of all sizes and types may be produced during the collision. Considering a IIB string theory in a (9+1)-dimensional space-time, interactions of Dirichlet (D) branes leads to the *unwinding* and subsequent *evaporation* of higher dimensionality branes, with the survival of three-dimensional (D3) branes embedded in a (9+1)-dimensional bulk — one of which could play the rôle of our universe — and D1-branes (D-strings) [5]. Large Fundamental (F) strings and D-strings that survive the cosmological evolution become cosmic superstrings. They are of cosmological size and could play the rôle of cosmic strings [6, 7], false vacuum remnants formed generically at the end of hybrid inflation within Grand Unified Theories [8, 9]. Cosmic superstrings have gained a lot of interest, particularly since it is believed that they may be observed in the sky, providing

both, a means of testing string theory and a hint for a physically motivated inflationary model (for a recent review, see e.g. Ref. [10]).

Brane collisions lead also to the formation of bound states, (p, q) -strings, which are composites of p F-strings and q D-strings [11, 12]. The presence of stable bound states implies the existence of junctions, where two different types of string meet at a point and form a bound state leading away from that point. Thus, when cosmic superstrings of different types collide, they can not intercommute, instead they exchange partners and form a Y -junction, as a consequence of charge conservation at the junction of colliding (p, q) -strings. The evolution of F, D strings and their bound states is a rather complicated problem, which necessitates both numerical as well as analytical investigations. Junctions may prevent the network from achieving a *scaling* solution, invalidating the cosmological model leading to their formation. In a number of studies, cosmic superstring evolution has been addressed via numerical experiments [13, 14, 15, 16, 17, 19]. The formation of three-string junctions and kinematic constraints for their collisions have been also investigated analytically [20, 21, 22].

In what follows, we address the question of the effect of junctions in the evolution of a cosmic superstring network, being composed by three components: p F-strings, q D-strings and their (p, q) bound states. The results of our numerical investigations can be summarised as follows. Firstly, there is clear evidence for *scaling* of all three components of the network, independently of the chosen initial configurations. Secondly, the existence of bound states effects the evolution of the network. Thirdly, for (p, q) strings there is a supplementary energy loss mechanism, in addition to the chopping off of loops, and it is this new mechanism that allows the network to scale. More precisely, the additional energy loss mechanism is the formation of bound states, whose length increases, lowering the overall energy of the network. We note that in our simulations we can have control over the initial population of bound states, and this renders our novel results particularly important.

2. The model

We adopt the model of Ref. [17]. This is a simple field theory model of (p, q) bound states, in analogy to the Abelian Higgs model, which incorporates the main features of string theory. To represent the two different species of strings, the model includes two complex scalar fields, ϕ and χ , of the Abelian Higgs model, coupled via a potential, so that bound states can be formed. The presence of stable bound states implies the existence of junctions, where two different types of string meet at a point and form a bound state leading away from that point.

We distinguish two different cases of cosmic superstring networks. The nature of cosmic superstrings depends on the particular brane inflationary model leading to their formation. In the case that both species of cosmic strings are BPS, the model is

described by the action [17]:

$$\mathcal{S} = \int d^3x dt \left[-\frac{1}{4}F^2 - \frac{1}{2}(D_\mu\phi)(D^\mu\phi)^* - \frac{\lambda_1}{4}(\phi\phi^* - \eta_1^2)^2 - \frac{1}{4}H^2 - \frac{1}{2}(D_\mu\chi)(D^\mu\chi)^* - \frac{\lambda_2}{4}\phi\phi^*(\chi\chi^* - \eta_2^2)^2 \right], \quad (1)$$

where the covariant derivative D_μ is defined by

$$\begin{aligned} D_\mu\phi &= \partial_\mu\phi - ie_1A_\mu\phi, \\ D_\mu\chi &= \partial_\mu\chi - ie_2C_\mu\chi. \end{aligned} \quad (2)$$

For clarity in our discussion, we label the ϕ field as “Higgs” and the χ field as “axion”, even though both fields are Higgs-like. The scalars are coupled to the $U(1)$ gauge fields A_μ and C_μ , with coupling constants e_1 and e_2 and field strength tensors $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ and $H_{\mu\nu} = \partial_\mu C_\nu - \partial_\nu C_\mu$, respectively. The scalar potentials are parametrised by the positive constants λ_1, η_1 and λ_2, η_2 , respectively.

In the case that one species of string is non-BPS, we remove the second gauge field by setting $e_2 = 0$. In this way, this species of string is represented by the topological defect of a complex scalar field with a global $U(1)$ symmetry. Such defects are characterised by the existence of long-range interactions [18] — as opposed to local strings in which all energy density is confined within the string, so that local strings have only gravitational interactions — implying different consequences for the evolution of the network.

In flat space the classical equations of motion for the ϕ and χ fields, obtained from the above action, Eq. (1), read [17]

$$\begin{aligned} \partial_\mu F^{\mu\nu} &= 2e_1 \text{Im}\phi^* D^\nu\phi, \\ \partial_\mu H^{\mu\nu} &= 2e_2 \text{Im}\chi^* D^\nu\chi, \\ D_\mu D^\mu\phi &= -2\lambda_1(\phi^*\phi - \eta_1^2)\phi - \lambda_2(\chi^*\chi - \eta_2^2)^2\phi \\ D_\mu D^\mu\chi &= -2\lambda_2\phi^*\phi(\chi^*\chi - \eta_2^2)\chi. \end{aligned} \quad (3)$$

By calculating numerically the tension of the (p, q) strings, for a range of values of the charges, and then fitting the data to the square-root expression

$$\mu_{(p,q)} = \mu_F \sqrt{p^2 + q^2/g_s^2}, \quad (4)$$

where μ_F denotes the effective fundamental string tension after compactification and g_s stands for the string coupling, we check [17] the validity of our model. We remind to the reader that Eq. (4) is exact only in the BPS limit.

The potential is chosen such that bound states are energetically preferred over single strings. This can be seen from the form of the potential for the axion strings. At the core of the Higgs string the field vanishes, $\phi = 0$, and therefore the potential of the axion vanishes as well. Therefore it is energetically favourable for the axion string to be located at the core of the Higgs string. In the true vacuum of the Higgs, we have $|\phi| = \eta_1$, so the axion string behaves just like the Higgs string.

We employ this model to study, via numerical simulations, overall properties of the cosmic superstring network. In particular, we are interested in investigating the effect of junctions and their rôle in achieving *scaling* of the network.

In the actual simulations we include the expansion of the universe by considering strings of fixed co-moving thickness, according to the model of Refs. [23, 24, 25]. In a Friedmann-Lemaître-Robertson-Walker universe and considering the temporal gauge, $A_0 = 0$, the equations of motion for the scalar fields take the form:

$$\partial_0 F_{0i} + 2(1-s) \frac{a'}{a} F_{0i} = 2a^{2s} e_1 \text{Im} \phi^* D^\nu \phi + \partial_j F_{ji} \quad (5)$$

$$\partial_0 C_{0i} + 2(1-s) \frac{a'}{a} C_{0i} = 2a^{2s} e_2 \text{Im} \chi^* D^\nu \chi + \partial_j C_{ji} \quad (6)$$

$$\phi'' + 2 \frac{a'}{a} \phi' + D_i D_i \phi = -a^{2s} \left[2\lambda_1 (\phi^* \phi - \eta_1^2) + \lambda_2 (\chi^* \chi - \eta_2^2)^2 \right] \phi \quad (7)$$

$$\chi'' + 2 \frac{a'}{a} \chi' + D_i D_i \chi = -2\lambda_2 a^{2s} \phi^* \phi (\chi^* \chi - \eta_2^2) \chi \quad (8)$$

where the time variable is the conformal time τ :

$$ds^2 = a^2(\tau) (-d\tau^2 + d\mathbf{x}^2) . \quad (9)$$

In our simulations we set $s = 0$, instead of the $s = 1$ required by the continuum-limit of the equations of motion (otherwise the defects will *fall through the lattice*); we also take, for simplicity, $a'/a = 1/\tau$.

We calculate the correlation length using the expression:

$$\xi = \sqrt{\frac{V}{L}} , \quad (10)$$

where V is the volume of the simulation box and L is the total length of the string network inside the box (following the same approach as for example in Ref. [19]). We identify the location of the string by the value of the scalar fields.

3. Results

We present our main results for the *scaling* of (p, q) string networks, using simulations of the classical time evolution on lattices with three spatial dimensions. We consider two types of networks, namely local-global and local-local networks. The axion field has a global $U(1)$ symmetry, and therefore the axion strings display long-range interactions.

The lattice discretisation of the equations of motion is discussed in Ref. [17]. We choose all parameters of the Lagrangian to have *natural* values,

$$\eta_1 = 1, \lambda_1 = 1, e_1 = 1, \eta_2 = 1, \lambda_2 = 1, e_2 = 1 . \quad (11)$$

The lattice spacing is $\delta x = 0.45$, roughly half the characteristic length scale set by the masses of the scalar and gauge fields in the broken phase. The time-step is $\delta t = 10^{-2}$. The simulations are carried out with boxes of volume 256^3 .

For each of the two (local-global and local-local) networks we consider two types of initial conditions: one in which a large percentage of strings (~ 50 % of the string

length) is in bound states, and the other one with a small percentage of strings ($\sim 2\%$ of the string length) is in bound states. We are able to achieve the small bound state percentage configuration, by starting with a large percentage configuration and then *shifting* one of the fields by roughly half the initial correlation length.

The type of behaviour we observe is that in most cases the string network does have a *scaling* behaviour, but the slope, γ , of the correlation length, ξ , versus time, defined by:

$$\xi(\tau) = \gamma\tau, \quad (12)$$

undergoes a discrete change as the network evolves. This change is most clearly visible in the case of a local-global string network, for both large and small amounts of bound states. The most dramatic change is that for the correlation length of the bound states themselves, provided the amount of bound states is small.

For local-local networks, the change in correlation length is not detectable if there is a large amount of bound states, while if there is a small amount of bound states we find a case where the bound states themselves do not exhibit *scaling* behaviour. This violation of *scaling* by the bound states can be explained by taking into account that the formation of bound states gives an additional possibility for a network of (p, q) strings to decrease its energy. Therefore, the length of the bound states increases as a function of time, violating the *scaling* behaviour which requires the total length to decrease. Once the length of the bound states have reached the maximum value, *scaling* resumes. This violation of *scaling* by the bound states does not lead to violation of *scaling* for the entire network, as it can be observed from the evolution of the Higgs and axion strings.

In Fig. 1 we show the string correlation length for the Higgs and axion fields, as well as for their bound states, as a function of time. The initial configuration is a local-global network with a large amount of bound states. Clearly, there is convincing evidence for scaling of the three components of the network. This scaling is characterised with a distinct change of the correlation length slope during the network evolution.




Figure 1. The Higgs (left), axion (middle), and bound state (right) string correlation length as a function of time. The network is a local-global one with a large amount of bound states. The data and linear fits for the two regimes are shown.

In Fig. 2 we show the string correlation length for the Higgs and axion fields, as well as for their bound states, as a function of time. The initial configuration is a

local-global network with a small amount of bound states. Again, scaling behaviour is apparent for all three components of the local-global network. The difference between the case plotted in Fig. 2 and the one plotted in Fig. 1, is the percentage of initial bound states. As one can easily observe from Fig. 2, if the initial state of a local-global network has a small amount of bound states, then the change of the correlation length slope during the evolution of the network is more acute.




Figure 2. The Higgs (left), axion (middle), and bound state (right) string correlation length as a function of time. The network is a local-global one with a small amount of bound states. The data and linear fits for the two regimes are shown.

We then draw the corresponding plots for local-local networks, with a large and a small amount of bound states in Fig. 3 and Fig. 4, respectively. We see, from Fig. 3, that for a local-local network with a large amount of bound states, all three components scale (as for the local-global case), whereas the change in correlation lengths during evolution is not really apparent (as opposed with the local-global case).




Figure 3. The Higgs (left), axion (middle), and bound state (right) string correlation length as a function of time. The network is a local-local one with a large amount of bound states. The data and linear fits for the two regimes are shown.

In Fig. 4, we plot the correlation length of the Higgs, the axion and their bound states for a local-local network with a small amount of bound states. Here we explicitly see that there is a violation of the *scaling* behaviour for the bound states, as a result of the increase of their length with time. As we have already mentioned earlier, this violation does not affect the entire network, which indeed scales.

In Fig. 5 we show, for comparison, the correlation length of a single species of string, using the same initial data as the one used for the (first) Higgs of the (p, q)




Figure 4. The Higgs (left), axion (middle), and bound state (right) string correlation length as a function of time. The network is a local-local one with a small amount of bound states. The data and linear fits for the two regimes are shown.

strings. Clearly, there is no change in the slope of the correlation length as a function of time. This is an important check to assure that finite size effects do not influence the results of the simulations.




Figure 5. The correlation length for a single species of cosmic string. There is no change in the slope of the correlation length as a function of time.

4. Conclusions

In the context of brane-world models, brane collisions lead to the formation of large Fundamental (F) strings and/or D1-branes (D-strings). Those who survive the cosmological evolution become cosmic superstrings, playing the rôle of their solitonic analogues. By observing strings in the sky, we may be able to test for the first (and maybe only) time string theory. To know the observational consequences of cosmic superstrings, it is essential to master the properties and evolution of such networks.

Performing numerical experiments, we have studied the dynamics and overall properties of superstring networks. More precisely, we have performed field theory simulations of p F- and q D-strings, and their (p, q) bound states. We have investigated the effect of bound states and in particular the approach to *scaling*, which is crucial for the cosmological consequences of cosmic superstring networks. Defects which do not scale are cosmologically undesired, since they may over-close the universe, leading to the *old* monopole problem.

Our studies have clearly shown that the three components of the network scale, independently of the chosen initial conditions. In addition, by having control over the initial abundance of bound states, we were able to identify the effect of bound states on the overall network evolution. Finally, we have found that there is an additional energy loss mechanism, beyond the chopping off loops. This new mechanism consists on the formation of bound states, resulting to a lower overall energy of the network, thus leading to scaling.

Acknowledgments

The work of M.S. is partially supported by the European Union through the Marie Curie Research and Training Network *UniverseNet* (MRTN-CT-2006-035863).

- [1] S. Sarangi and S. H. H. Tye, Phys. Lett. B **536**, 185 (2002) [arXiv:hep-th/0204074];
- [2] N. T. Jones, H. Stoica and S. H. H. Tye, Phys. Lett. B **563**, 6 (2003) [arXiv:hep-th/0303269];
- [3] G. Dvali and A. Vilenkin, J. Cosm. & Astrop. Phys. **0403** (2004) 010 [arXiv:hep-th/0312007].
- [4] G. Dvali and S. H. Tye, Phys. Lett. B **450** (1999) 72 [arXiv:hep-ph/9812483].
- [5] R. Durrer, M. Kunz and M. Sakellariadou, Phys. Lett. B **614**, 125 (2005) [arXiv:hep-th/0501163].
- [6] A. Vilenkin and E. P. S. Shellard, *Cosmic Strings and Other Topological Defects* (Cambridge University Press, Cambridge, 1994).
- [7] M. Sakellariadou, Springer Lecture Notes in Physics, **718** (2007) 247 [arXiv:hep-th/0602276].
- [8] R. Jeannerot, J., Rocher and M. Sakellariadou, Phys. Rev. D **68** (2003) 103514 [arXiv:hep-ph/0308134].
- [9] M. Sakellariadou, Springer Lecture Notes in Physics, **738** (2008) 359 [arXiv:hep-th/0702003].
- [10] M. Sakellariadou, *Cosmic Superstrings*, Philosophical Transactions A, Royal Society (to appear) [arXiv:0802.3379 [hep-th]].
- [11] E. Copeland, R. C. Myers, L. Polchinski, J. of High Energy Phys. **0406** (2004) 013.
- [12] L. Leblond, S. H. Tye, J. High Energy Phys. **0403** (2005) 055.
- [13] M. Sakellariadou, JCAP **0504** (2005) 003 [arXiv:hep-th/0410234].
- [14] A. Avgoustidis and E. P. S. Shellard, Phys. Rev. D **71** (2005) 123513 [arXiv:hep-ph/0410349].
- [15] E. J. Copeland and P. M. Saffin, JHEP **0511** (2005) 023 [arXiv:hep-th/0505110].
- [16] M. Hindmarsh and P. M. Saffin, JHEP **0608** (2006) 066 [arXiv:hep-th/0605014].
- [17] A. Rajantie, M. Sakellariadou and H. Stoica, JCAP **0711** (2007) 021 [arXiv:0706.3662 [hep-th]].
- [18] G. D. Moore and H. Stoica, Phys. Rev. D **74**, 065003 (2006) [arXiv:hep-th/0605070].
- [19] J. Urrestilla and A. Vilenkin, JHEP **0802** (2008) 037 [arXiv:0712.1146 [hep-th]].
- [20] E. J. Copeland, T. W. B. Kibble and D. A. Steer, Phys. Rev. Lett. **97** (2006) 021602.
- [21] E. J. Copeland, T. W. B. Kibble and D. A. Steer, Phys. Rev. D **75** (2007) 065024 [arXiv:hep-th/0611243].
- [22] E. J. Copeland, H. Firouzjahi, T. W. B. Kibble and D. A. Steer, Phys. Rev. D **77** (2008) 063521 [arXiv:0712.0808 [hep-th]].
- [23] B. S. Ryden, W. H. Press and D. N. Spergel “The evolution of networks of domain walls and cosmic strings”, CFA-3011 (1989).
- [24] W. H. Press, B. S. Ryden and D. N. Spergel “Dynamical evolution of domain walls in an expanding universe”, NSF-ITP-89-51, CFA-1870 (1989).
- [25] J. N. Moore, E. P. S. Shellard and C. J. A. Martins, Phys. Rev. D **65**, 023503 (2002) [arXiv:hep-ph/0107171].